

Comment on “Phase transition for quenched coupled replicas in a plaquette spin model of glasses”

Giulio Biroli*

*IPhT, CEA/DSM-CNRS/URA 2306, CEA Saclay, F-91191 Gif-sur-Yvette Cedex, France
LPS, Ecole Normale Supérieure, 24 rue Lhomond, 75231 Paris Cedex 05 - France*

Gilles Tarjus[†] and Marco Tarzia[‡]

*LPTMC, CNRS-UMR 7600, Université Pierre et Marie Curie,
boîte 121, 4 Pl. Jussieu, 75252 Paris cédex 05, France*

(Dated: December 16, 2015)

PACS numbers:

In a recent letter,¹ Jack and Garrahan have studied the thermodynamic phase transition associated with an overlap order parameter in spin models of glasses (PSMs) formed by plaquettes of p spins. The glassy features of such models are known to be well described in terms of rare defects and kinetic constraints. Yet, the authors give strong evidence that the 3-dimensional square pyramid model (SPyM) displays a transition line in the temperature (T)-external field (ϵ) plane which connects a zero-temperature critical point at $T_c(0) = 0$ to a finite-temperature one at $T_c(\epsilon_c) > 0$. This result, which complements an earlier study by the same authors² on the “annealed” calculation for two coupled replicas in $d = 2$ (triangular plaquette model or TPM) and in $d = 3$ (SPyM), shows that the PSMs behave similarly to what expected from the random first-order transition (RFOT) theory, based on the mean-field (MF) theory of the glass transition, with the notable exception that the Kauzmann temperature $T_K = T_c(0)$ is equal to zero.

Here we show that the properly implemented MF approach for PSMs turns out to be surprisingly predictive for the finite-dimensional behavior. As displayed in Fig. 1, the MF phase diagram of the SPyM on a Bethe lattice with the same connectivity, $c = p = 5$, as in $d = 3$ is remarkably close to its $d = 3$ counterpart in the annealed and the quenched settings.^{1,2} (It is obtained through standard techniques based on recursion relations and the cavity method.⁴⁻⁷) Both the singular behavior of $T_c(\epsilon)$ and the result $T_K = T_c(0) = 0$ are well captured.⁸ Only, as expected, the MF line terminates at higher T_c and ϵ_c . At odds with naive anticipations, the fact that $T_K = 0$ is *not* the result of long-distance fluctuations but comes from the mere local-connectivity constraint $c = p$: indeed, for $c > p$, MF predicts $T_K > 0$.^{5,6} Note that very poor results (in particular $T_K > 0$) would have instead been obtained by using a MF approach based on completely connected lattices. As discussed in Ref. [3], the RFOT found in infinite d is very fragile to short-ranged fluctuations associated with local properties, such as the connectivity of the underlying lattice for spin models. It is therefore crucial to focus on a MF description, such as the Bethe approximation, that takes them into account.³

All in all, and contrary to expectations, the MF de-

scription appears robust with respect to fluctuations as far as overlap properties go.⁹ Finite- d fluctuations enforce convexity of the potential $V(q)$ (the free-energy cost for maintaining an overlap q with a reference configuration,¹⁰ sketched in Fig. 1) and prevent true metastability. Nonetheless, the physics of the unbiased system in $\epsilon = 0$ is well described by MF up to a scale, the point-to-set length ξ_{PTS} , that diverges as $T \rightarrow 0$;¹¹ $V(q)$ remains singular in $d = 3$ (with a linear segment) and a configurational entropy s_c can then be univocally defined at low enough T , even in finite d (see Fig. 1). Although one expects $s_c \sim e^{-1/T}$ as $T \rightarrow T_K = 0$ in both MF and $d = 3$,¹ other critical exponents are likely to be different, as, *e.g.*, that characterizing the relation between ξ_{PTS} and s_c . Given the robustness of the thermodynamic MF predictions, understanding the implications for the dynamics and the connection to the defect picture is a key question.¹² Further studies of the PSMs could be instrumental in clarifying it.




FIG. 1: Phase diagram of the SPyM in both the annealed and the quenched descriptions for the Bethe lattice with $c = 5$ (dashed lines) and for $d = 3$ (full lines^{1,2}). Inset: Sketch of the potential $V(q)$ in MF (left) and for $d = 3$ (right).

-
- * Electronic address: giulio.biroli@cea.fr
† Electronic address: tarjus@lptmc.jussieu.fr
‡ Electronic address: tarziah@lptmc.jussieu.fr
- ¹ R. L. Jack and J. P. Garrahan, arXiv:1508.06470 (2015).
 - ² R. M. Turner, R. L. Jack, and J. P. Garrahan, Phys. Rev. E **92**, 022115(2015).
 - ³ C. Cammarota *et al.*, Phys. Rev. B **87**, 064202 (2013).
 - ⁴ M. Mézard and G. Parisi, Europhys. J. B **20**, 217 (2001).
 - ⁵ L. Zdeborová and F. Krzakala, Phys. Rev. B **81**, 224205 (2010); Y. Matsuda *et al.*, J. Phys. A: Math. Theor. **44**, 185002 (2011).
 - ⁶ S. Franz, G. Gradenigo, and S. Spigler, arXiv:1507.0572 (2015).
 - ⁷ G. Biroli, G. Tarjus, and M. Tarzia, in preparation (2015).
 - ⁸ In the annealed case, the self-dual property identically applies to $d = 3$ and to the MF with $c = p = 5$.
 - ⁹ In $d = 2$, long-range fluctuations in the quenched setting do suppress the transition line otherwise present for $\epsilon = 0$ in the MF TPM with $c = p = 3$.
 - ¹⁰ S. Franz and G. Parisi, J. Phys. (Paris) I **5**, 1401 (1995).
 - ¹¹ R. L. Jack and J. P. Garrahan, J. Chem. Phys. **123**, 164508 (2005).
 - ¹² See also L. Foini, F. Krzakala, and F. Zamponi, JSTAT, P06013 (2012).